



पुर्णा International School

Shree Swaminarayan Gurukul, Zundal

Class – XII

Subject: Chemistry(Practical)

Experiment (2021_22)

Exp. No	Aim
	QUANTITATIVE ANALYSIS
1	Prepare 250 ml of 0.1M Solution of Oxalic Acid From Crystalline Oxalic Acid
2	Determination of Concentration/Molarity of KMnO_4 Solution by Titrating it against a 0.1M Standard Solution of Oxalic acid
3	Determination of Concentration/Molarity of KMnO_4 Solution by Titrating it against a Standard Solution of Ferrous ammonium sulphate
	QUALITATIVE ANALYSIS
4	To Identify the given inorganic salt $[\text{Ba}(\text{NO}_3)_2]$
5	To Identify the given inorganic salt $[\text{Pb}(\text{CH}_3\text{COO})_2]$
6	To Identify the given inorganic salt $[\text{Pb}(\text{NO}_3)_2]$
7	To Identify the given inorganic salt PbCl_2
8	To Identify the given inorganic salt MgSO_4
9	To Identify the given inorganic salt ZnCl_2
10	To Identify the given inorganic salt ZnSO_4
11	To Identify the given inorganic salt $[(\text{NH}_4)_3\text{PO}_4]$
12	To Identify the given inorganic salt NH_4Br
13	To Identify the given inorganic salt $[(\text{NH}_4)_2\text{CO}_3]$
14	To Identify the given inorganic salt $[\text{Sr}(\text{NO}_3)_2]$

	ORGANIC COMPOUNDS
15	To Identify functional group of Aldehyde (-CHO)
16	To Identify functional group of Ketone (-CO-)
17	To Identify functional group of Alcohol (-OH)
18	To Identify functional group of Carboxylic acid (-COOH)

EXPERIMENT 11.1



Prepare 250 ml of 0.1 M solution of oxalic acid from crystalline oxalic acid.

THEORY

Molecular mass of crystalline oxalic acid $\left(\begin{array}{c} \text{COOH} \\ | \\ \text{COOH} \end{array} \cdot 2\text{H}_2\text{O} \right) = 126$

Hence, for preparing 1000 ml of 1M oxalic acid, weight of oxalic acid crystals required = 126 g

∴ For preparing 250 ml of 0.1M solution,

$$\text{oxalic acid crystals required} = \frac{126}{1000} \times 250 \times 0.1 = 3.150 \text{ g.}$$

APPARATUS

Watch glass, analytical balance, weight box, fractional weight box, 250 ml beaker, glass rod, 250 ml measuring flask and wash bottle.

CHEMICALS REQUIRED

Oxalic acid crystals and distilled water.

PROCEDURE

1. Take a watch glass, wash it with distilled water and then dry it.
2. Weigh the clean and dried watch glass accurately and record its weight in the notebook.
3. Weigh 3.150 g oxalic acid on the watch glass accurately and record this weight in the notebook.
4. Transfer gently and carefully the oxalic acid from the watch glass into a clean 250 ml measuring flask using a funnel. Wash the watch glass with distilled water with the help of a wash bottle to transfer the particles sticking to it into the funnel [Fig. 11.14]. The volume of distilled water for this purpose should not be more than 50 ml.
5. Finally wash the funnel well with distilled water with the help of a wash bottle to transfer the solution sticking to the funnel into the measuring flask [Fig. 11.15].
6. Swirl the measuring flask till solid oxalic acid dissolves.
7. Add enough distilled water to the measuring flask carefully, upto just below the etched mark on it, with the help of a wash bottle.




Fig. 11.14. Transferring oxalic acid to the flask.




Fig. 11.15. Adding water.

8. Add the last few drops of distilled water with a pipette or a dropper until the lower level of the meniscus just touches the mark on the measuring flask [Fig. 11.16].
9. Stopper the measuring flask and shake gently to make the solution uniform throughout. Label it as 0.1 M oxalic acid solution [Fig. 11.17].




Fig. 11.16. Adding last small amount of water dropwise.




Fig. 11.17. Standard solution of oxalic acid.


EXPERIMENT 6.1

Aim



To determine the concentration/molarity of KMnO_4 solution by titrating it against a 0.1 M standard solution of oxalic acid.

Reactions of oxalic acid



A. Chemical equations



B. Ionic equation



Material Required



	• Measuring flask (250 mL):	One		• Oxalic acid	: As per need
	• Burette (50 mL)	: One		• Potassium permanganate solution	: As per need
	• Burette stand	: One		• 1.0 M Sulphuric acid	: As per need
	• Pipette	: One			
	• Conical flask	: One			
	• Funnel	: One			
	• Weighing bottle	: One			
	• Glazed tile(white)	: One			
	• Burner	: One			
	• Wire gauze	: One			
	• Chemical balance	: One			

Procedure

A. Preparation of 0.1 M standard solution of oxalic acid

Prepare 0.1M oxalic acid solution as mentioned in experiment 2.1 (Unit 2, Class XI, Laboratory Manual)

B. Titration of oxalic acid solution against potassium permanganate solution

Oxalic acid	
Potassium permanganate	 
Sulphuric acid	

- Rinse and fill a clean burette with potassium permanganate solution. Remove the air bubble, if any, from the nozzle of the burette by releasing some solution through it. The burette used in the permanganate titration must have a glass stop cock as rubber is attacked by permanganate ions.
- Take 10 mL of 0.1 M oxalic acid solution in a conical flask and add half of the test tube full (= 5 mL) of 1.0 M H_2SO_4 to it to prevent the formation of any precipitate of manganese dioxide during the course of the titration.
- Heat the oxalic acid solution upto $50^\circ - 60^\circ\text{C}$ before titrating it with potassium permanganate solution taken in the burette. To increase the visibility of the colour change, place the conical flask containing the solution to be titrated over a white glazed tile kept below the nozzle of the vertically fitted burette.
- Note the initial reading of the volume of permanganate solution in the burette and add it in small volumes to the hot oxalic acid solution while swirling the contents of the flask gently. The violet colour of permanganate solution is

discharged on reaction with oxalic acid. The end point is indicated by the appearance of permanent light pink colour due to a slight excess of permanganate solution.

- (v) Repeat the titration till three concordant readings are obtained. Since the solution of KMnO_4 is of dark colour, the upper meniscus should be considered for noting the burette readings.
- (vi) Record the readings as shown in observation Table 6.1 and calculate the strength of potassium permanganate solution in mols/litre.

Table 6.1 : Titration of potassium permanganate solution against standard oxalic acid solution

Sl. No.	Volume of Oxalic acid in mL	Burette readings		Volume (V) of KMnO_4 used $V = (y-x)$ mL
		Initial (x)	Final (y)	

6/10

Calculations

- (i) The strength of the unknown solution in terms of molarity may be determined by the following equation.

$$a_1 M_1 V_1 = a_2 M_2 V_2 \quad (6.1)$$

For oxalic acid vs potassium permanganate titration:


$a_1 = 2$. (the number of electrons lost per formula unit of oxalic acid in a balanced equation of half cell reaction)

$a_2 = 5$. (the number of electrons gained per formula unit of potassium permanganate in the balanced equation of half cell reaction)

M_1 and M_2 are the molarities of oxalic acid and potassium permanganate solutions used in the titration.

V_1 and V_2 are the volumes of oxalic acid and potassium permanganate solutions.

On putting the value of a_1 and a_2 in equation 6.1 we get



$$M_2 = \frac{2M_1V_1}{5V_2} \quad (6.2)$$

We can calculate the molarity of potassium permanganate solution by using equation 6.2. Strength of the solution is given by the following equation:

$$\text{Strength} = \text{Molarity} \times \text{Molar mass}$$

Result

- (i) Molarity of KMnO_4 solution is _____.
- (ii) Strength of KMnO_4 solution is _____.

Precautions

- (a) Always rinse the burette and the pipette with the solutions to be taken in them.
- (b) Never rinse the conical flask with the experimental solutions.
- (c) Remove the air gaps if any, from the burette.
- (d) Never forget to remove the funnel from the burette before noting the initial reading of the burette.
- (e) No drop of the liquid should hang at the tip of the burette at the end point and while noting reading.
- (f) Always read the upper meniscus for recording the burette reading in the case of all coloured solutions.
- (g) Never use pipette and burette with a broken nozzle.
- (h) Lower end of the pipette should always remain dipped in the liquid while sucking the liquid.
- (i) Do not blow out the last drop of the solution from the jet end of the pipette.
- (j) The strength of the solution must be calculated up to the fourth decimal place.
- (k) Do not forget to heat the mixture of oxalic acid and H_2SO_4 solutions between 50°C and 60°C while titrating it against potassium permanganate.

7/10

EXPERIMENT 6.2




Aim

To determine the concentration/molarity of KMnO_4 solution by titrating it against standard solution of ferrous ammonium sulphate.




Theory

Like oxalic acid, ferrous ammonium sulphate also acts as a reducing agent in the titration against potassium permanganate. The reaction which takes place is given below :



(a) Chemical equation



(b) Ionic equation



Material Required

	• Measuring flask (250 mL) :	One		• Potassium permanganate solution :	As per need
	• Burette (50 mL) :	One		• Dilute sulphuric acid :	As per need
	• Burette stand :	One		• Ferrous ammonium sulphate :	As per need
	• Pipette :	One			
	• Conical flask :	One			
	• Glazed tile (white) :	One			
	• Funnel :	One			
	• Weighing bottle :	One			

Procedure

Potassium permanganate



Sulphuric acid

**A. Preparation of 0.05 M, standard solution of ferrous ammonium sulphate**

(Molar mass of $\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O} = 392 \text{ g mol}^{-1}$).

- (i) Weigh 4.9000 g of ferrous ammonium sulphate and transfer it into a 250 mL measuring flask through a funnel.
- (ii) Transfer the solid sticking to the funnel with the help of distilled water into the flask and add dilute H_2SO_4 into the flask drop wise to get the clear solution.
- (iii) Shake the flask till the substance dissolves and make the solution upto the mark.

B. Titration of ferrous ammonium sulphate against potassium permanganate solution

- (i) Rinse and fill the clean burette with potassium permanganate solution. Remove air bubbles if any, from the burette tip by releasing some solution through it.
- (ii) Take 10 mL of 0.05 M ferrous ammonium sulphate solution in a conical flask and add half test tube ($\approx 5 \text{ mL}$) full of (1.0 M) H_2SO_4 to it.
- (iii) Titrate the above solution with potassium permanganate solution till the colour of the solution changes to permanent pink. Swirl the content of the flask during the titration.
- (iv) Repeat the titration, until three concordant readings are obtained.
- (v) Record the readings as shown in observation Table 6.2 and calculate the strength of potassium permanganate solution in mols/litre.

Table 6.2 : Titration of potassium permanganate solution against standard ferrous ammonium sulphate solution

Sl. No.	Volume of ferrous ammonium sulphate solution used for each titration in mL	Burette Readings		Volume (V) of KMnO_4 used $V = (y-x)$ mL
		Initial (x)	Final (y)	

Calculations

The strength of unknown solution in terms of molarity may be determined by the following equation :

$$a_1 M_1 V_1 = a_2 M_2 V_2$$

M_1 and M_2 are the molarities of ferrous ammonium sulphate and potassium permanganate solutions and V_1 and V_2 are volumes of ferrous ammonium sulphate and potassium permanganate solutions, respectively.

$a_1 = 1$, (the number of electrons lost per formula unit of ferrous ammonium sulphate in the half cell reaction)

$a_2 = 5$, (the number of electrons gained per formula unit of potassium permanganate in a half cell reaction)

Strength can be calculated by the formula given below :

$$\text{Strength} = \text{Molarity} \times \text{Molar mass}$$

Result

The strength of the given potassium permanganate solution is ____ g/L.

Precautions

- Always use a fresh sample of ferrous ammonium sulphate to prepare its standard solution.
- Other precautions are same as that in Experiment 6.1.

AIM

EXPERIMENT-5

AIM - To identify the given inorganic salt $[Ba(NO_3)_2]$

EXPERIMENT	OBSERVATIONS	INFERENCE
* PRIMARY TEST		
1. Color.	white	Absence of Cu^{2+} , Fe^{3+} , Co^{+2} , Mn^{+2}
2. Smell	No specific	NH_4^+ , S^{2-} , CH_3COO^- absent.
3. Gas evolved	A reddish brown gas evolved, which turned $FeSO_4$ sol ⁿ black.	NO_3^- may be present.
4. Sublimation	No sublimation	NH_4^+ , I^- may be absent.
5. Decrepitation	No decrepitation	$[Pb(NO_3)_2]$, $NaCl$, KBr absent.
6. Residue	white	Zn^{2+} , Pb^{2+} may be absent.
7. FLAME TEST - Prepare a paste Salt + Conc. HCl	Persistent grassy green flame on prolonged heating.	Ba^{+2} may be present.
8. Salt + dil. H_2SO_4 (warm)	No gas evolves	CO_3^{2-} , S^{2-} , NO_2 may be absent.
9. Salt + dil. H_2SO_4 adding drops of $KMnO_4$	Pink color of $KMnO_4$ is lost.	Cl^- , Br^- , I^- , $C_2O_4^{2-}$, Fe^{2+} may be absent.
10. Heat a pinch of salt and conc. NaOH	No ammonia gas evolved	NH_4^+ absent.

AIM:			
11.	Preparation of (0.5) shake a salt + water mix	sol ^m obtained	label it as the original solution.
12.	To a part of (0.5) add 1-2 ml of dil. HCl	No ppt formed	GI Absent (Pb ²⁺ absent).
13.	To a part of sol ^m pass H ₂ S gas.	No ppt formed	GII Absent. (Pb ²⁺ , Cu ²⁺ , Ag ³⁺ absent)
14.	To remaining sol ^m . add solid NH ₄ Cl, Boil, cool down, add a few drops NH ₄ OH	No ppt formed	GIII absent. (Fe ²⁺ , Al ³⁺ , absent)
15.	Through a part of this sol ^m , pass H ₂ S gas.	No ppt formed	GIV absent. (Zn ²⁺ , Mn ²⁺ , Ni ²⁺ , Co ²⁺ absent).
16.	To the remaining ammonical solution, add ammonium carbonate.	white ppt formed	GIV present (Ca ²⁺ , Ba ²⁺ , Sr ²⁺ may be present).
17.	Co		
*	CONFIRMATORY TEST		
17	For Nitrate -		
a)	Copper chips test, heated a pinch of the salt with conc. H ₂ SO ₄	Reddish Brown Gas	NO ₃ ⁻ Confirmed

AIM:			
b)	Ring Test	Dark Brown ring is observed	NO_3^- Confirmed.
18.	For Ba^{2+} -		
a)	Potassium Chromate test	Yellow ppt	Ba^{2+} Confirmed
b)	Perform flame test with salt	Greenish green flame	Ba^{2+} Confirmed

RESULT -

- i) Acid Radical - NO_3^-
- ii) Basic Radical - Ba^{2+}

PRECAUTIONS -

- i) Handle reagents properly
- ii) Never heat a wet test-tube
- iii) Keep processing test-tube away from body.
- iv) Don't inhale unknown / poisonous gases.

AIM:

EXPERIMENT-6

AIM - To identify the given inorganic salt $[Pb(CH_3COO)_2]$

EXPERIMENT	OBSERVATION	INFERENCE
* PRIMARY TEST-		
1. Colour	white	Shows absence of $Cu^{2+}, Ni^{2+}, Fe^{3+}, Mn^{2+}, Co^{2+}$.
2. Smell	Vinegar like smell	Shows presence of CH_3COO^-
3. Density	heavy	Salt of Pb^{2+} or Ba^{2+} carbonate may be present.
4. Deliquescence	No deliquescence	Shows absence of $Zn(NO_3)_2$ & Cl of Zn^{+2}, Mg^{+2} etc.

* DRY HEATING TEST

5. Gas evolved	Colourless gas with characteristic vinegar like smell	CH_3COO^- may be present.
6. Sublimate formed	No sublimation	NH_4^+ & I^- are absent.
7. Descr. precipitation	No descr. precipitation	Salts like $Pb(NO_3)_2, NaCl, KI$ are absent.
8. Residue.	White salt becomes black on heating	CH_3COO^- may be present.

* FLAME TEST-

9. Make a paste of salt and conc. HCl & perform the test	Dull bluish white flame	Pb^{2+} may be present.
--	-------------------------	---------------------------

VISION

Teacher's Signature : _____

AIM:			
10.	Dil H_2SO_4 test: Treat a pinch of salt with dil. H_2SO_4 (+Heat)	No gas evolved.	CO_3^{2-} , S^{2-} , NO_2^- , SO_3^{2-} may be absent.
11.	KMnO ₄ Test: To a pinch of salt add dil H_2SO_4 & heat then add KMnO ₄	Pink color of KMnO ₄ wasn't discharged	Cl^- , Br^- , I^- , $C_2O_4^{2-}$, Fe^{2+} may be absent.
12.	Heat a pinch of salt with conc. NaOH	No ammonia gas is evolved	NH_4^+ absent.
13.	Preparation of (o.s.): Shake mix of salt & water	Solution is obtained	Label it as the original solution.
14.	To a part of o.s. add 2ml of dil HCl	white ppt is obtained	<ul style="list-style-type: none"> • Group I is present • Pb^{2+} might be present.

RESULT - i) Acid Radical : CH_3COO^-
 ii) Basic Radical : Pb^{2+}

PRECAUTIONS - i) Don't heat wet test tube
 ii) Don't inhale gases, they might be poisonous.
 iii) Keep test-tube far from face, while dry heating.
 iv) Handle reagents carefully.

AIM:

EXPERIMENT - 7.

AIM - To identify the given inorganic salt $Pb(NO_3)_2$

EXPERIMENT	OBSERVATION	INFERENCE
* PRIMARY TEST -		
1. Color.	white.	Shows absence of NH_4^+ , CH_3COO^- , S^{2-}
2. Smell	No specific odour	Shows absence of Cu^{2+} , Ni^{2+} , Fe^{2+} , Co^{2+} .
3. Density.	Heavy / Thick.	Salt of Pb^{2+} or Ba^{2+} carbonate.
4. Deliquescence.	No deliquescence.	Shows absence of $Zn(NO_3)_2$, Chlorides of Zn^{2+} , Mg^{2+} etc.
* DRY HEATING TEST -		
5. Gas evolved.	A reddish brown gas evolved which turned $FeSO_4$ solution, black.	NO_3^- may be present.
6. Sublimate formed	No Sublimation.	Shows absence of NH_4^+ & I^-
7. Descrripitation.	The salt descrripitates	$Pb(NO_3)_2$, $NaCl$, KBr may be present.
8. Swelling.	No Swelling.	Shows absence of indicated. PO_4^{3-}
9. Residue	Hot \rightarrow Brown. Cold \rightarrow Yellow.	Pb^{2+} might be present.

AIM:

FLAME TEST

10.	Prepare a paste of Salt 9 with concentrated HCl and perform flame test.	Dull bluish - white flame	Pb^{2+} Pb^{2+} may be present.
11.	Dil. H_2SO_4 test: Treat a pinch of salt with dil. H_2SO_4 & heat.	No gas evolved	Pb^{2+} may be present.
12.	$KMnO_4$ Test: To a pinch of salt add dil. H_2SO_4 (hot) and then add a drop of $KMnO_4$.	Pink color of $KMnO_4$ was not discharged.	Cl^- , Br^- , I^- , $C_2O_4^{2-}$, Fe^{2+} may be absent.
13.	Conc. H_2SO_4 Test: Salt + Conc. H_2SO_4 + (Heat if required)	A reddish brown gas evolved, which turned $FeSO_4$ sol ⁿ into black.	NO_3^- may be present.
14.	Confirmatory Test: FOR NITRATE -		
a)	'Cu' chip test; Heat a small quantity of salt with conc. H_2SO_4 and a few 'Cu' chips	Reddish brown gas evolves	NO_3^- is confirmed $2KNO_3 + H_2SO_4 \rightarrow K_2SO_4 + 2HNO_3$
b)	Ring Test - 2-3ml of salt sol ⁿ + $FeSO_4$ sol ⁿ Add conc. H_2SO_4 along sides of test tube.	Solution obtained of dark brown color, 2 liquids at the junction.	label it as original sol ⁿ .
15.	Preparation of (0.5) Shake a pinch of salt with water.	Solution obtained	label it as original sol ⁿ .

AIM:

16.	To a pinch of O.S., add 1-2 ml of dil. HCl	white ppt is formed	Group I: Pb^{2+} may be present
17.	Confirmatory test for Pb^{2+} dissolve white ppt with distilled water & divide it in two parts.		
a)	KI Test: To one part add KI solution.	Yellow ppt obtained	Pb^{2+} is confirmed $PbCl_2 + 2KI \rightarrow PbI_2 + 2KCl$
b)	K_2CrO_4 Test: to one part add K_2CrO_4 sol ⁿ .	Yellow ppt obtained	Pb^{2+} is confirmed $PbCl_2 + K_2CrO_4 \rightarrow PbCrO_4 + 2KCl$

RESULT-

Acid Radical: NO_3^- Basic Radical: Pb^{2+} .

PRECAUTIONS -

- 1) Don't heat a wet test tube.
- 2) Don't inhale gases directly, they might be harmful.
- 3) Keep test tube away from face while dry heating.
- 4) Handle reagents carefully.

AIM:

EXPERIMENT-8

AIM- To identify the given inorganic salt $Pb(NO_3)_2$

EXPERIMENT	OBSERVATION	INFERENCE
* PRIMARY TEST-		
1. Colour	white	Shows absence of Cr^{2+} , Ni^{2+} , Fe^{2+} , Fe^{3+} , Mn^{2+} , Co^{2+}
2. Smell	No specific smell	Shows absence of NH_4^+ , CH_3COO^- .
3. Density	Heavy	Salt of Pb^{2+} or Ba^{2+} Carbonate may be present.
4. Deliquescence.	no deliquescence	Shows absence of $Zn(NO_3)_2$ & chlorides of Zn^{2+} .
* DRY HEATING TEST		
5. Gas evolved	Colourless gas with pungent smell, white fumes with ammonia \rightarrow white ppt with $AgNO_3$.	Cl^- may be present.
6. Description.	No description.	$Pb(NO_3)_2$, $NaCl$ are absent
7. Residue.	Hot \rightarrow Brown. Cold \rightarrow Yellow	Pb^{2+} might be present.

AIM:

* FLAME TEST -

8.	Make a paste of Salt + conc. HCl.	Dull bluish white flame.	Pb^{2+} may be present.
9.	Dil. H_2SO_4 Test. Treat a pinch of Salt with dil H_2SO_4 and heat.	No gas evolved	CO_3^{2-} , S^{2-} , NO_2^- , SO_3^{2-} may be present.
10.	Conc. H_2SO_4 + Salt + Heat (if required)	Colourless gas, pungent smell, white fumes with ammonia, and white $AgNO_3$ ppt.	Cl^- may be present.
11.	Heat a pinch of Salt with conc. NaOH	No ammoniac gas evolved	NH_4^+ absent.
12.	Preparation (os) of: Shaking a mix of Salt and water	Solution obtained	Label it as the original Sol ⁿ .
13.	To a part of 0.5 add 1.2 ml of dil. HCl.	white colored ppt is obtained	Group I is present Pb^{2+} may be present.
14.	Heat a pinch of Salt with conc. NaOH.	No NH_3 gas is released	Absence of NH_4^+ ions.

RESULT -

Acid Radical - Cl^-

Basic Radical - Pb^{2+}

AIM

PRECAUTIONS -

- i) Never heat wet test-tube
- ii) No direct inhalation of gases.
- iii) Keep tube away from face while dry heating.
- iv) Handle reagents carefully.

AIM:

EXPERIMENT-9

AIM - To identify the given inorganic salt $MgSO_4$

EXPERIMENT	OBSERVATIONS	INFERENCE
* PRIMARY TEST-		
1. Colour	White	Shows absence of Cu^{2+} , Ni^{2+} , Mn^{2+} , Co^{2+} , NH_4^+ , CH_3COO^- , S^{2-}
2. Smell	Odourless	S^{2-} , NH_4^+ , CH_3COO^- are absent.
3. Gas evolved	No gas evolved	S^{2-} , SO_3^{2-} , Cl^- , CH_3COO^- , NH_4^+ , NO_3^- are absent.
4. Sublimate formed	No sublimation	NH_4^+ , I^- are absent.
5. Decipitation	No decipitation.	$Pb(NO_3)_2$, $NaCl$, KBr , are absent.
6. Residue	White residue that glows on heating	Ba^{2+} , Sr^{2+} , Ca^{2+} , Hg^{2+} , Al^{3+} maybe present.
7. Flame Test Make a paste of salt + conc. HCl	No specific flame color	Ca^{2+} , Sr^{2+} , Ba^{2+} , Cu^{2+} , Zn^{2+} , Pb^{2+} may be absent.
8. Dil H_2SO_4 Test Treat a pinch of salt + dil H_2SO_4 and heat	No gas evolved	CO_3^{2-} , S^{2-} , NO_2^- , SO_3^{2-} might be absent.
9. $KMnO_4$ Test: A pinch of salt added to dil. H_2SO_4 & heat.	Pink color of $KMnO_4$ wasn't discharged.	Cl^- , Br^- , I^- , $C_2O_4^{2-}$, Fe^{2+} maybe absent.

AIM	Observation	Inference
10. Then add $KMnO_4$. Conc. H_2SO_4 + Salt + Heat (if required)	No gas evolved	Cl^- , Br^- , I^- , NO_3^- , (CH_3COO^-) are absent.
11. Heat a pinch of Salt with conc. $NaOH$	No ammonia gas evolved	NH_4^+ absent.
12. Shake a mix of Salt with water.	Solution obtained	Label as original solution.
13. To a part of 0.5 add 1-2 ml of dil. HCl	No ppt formed	Group I absent Pb^{2+} absent.
14. Through the above formed solution, Pass H_2S gas	No ppt formed	Group II absent (Pb^{2+} , Cu^{2+} , As^{2+} , etc.)
15. To remaining Sol^m add a pinch of solid NH_4Cl , Boil the Sol^m and add excess NH_4OH	No ppt formed.	Group III absent (Fe^{2+} , Al^{3+} , absent)
16. To the remaining Sol^m add ammonium carbonate.	No ppt formed	Group V absent (Ca^{2+} , Ba^{2+} absent).
17. Through a part of the above Sol^m , pass H_2S gas	No ppt formed	Group IV absent (Zn^{2+} , Mn^{2+} , Ni^{2+} , Co^{2+} absent)

RESULT -

Acid Radical - SO_4^{2-}

Basic Radical - Mg^{2+}

Expt. No. _____

AIM:

PRECAUTIONS

- i) Don't heat wet test-tube
- ii) Don't inhale any gases.
- iii) Keep tube away from face while heating dry.
- iv) Handle reagents carefully.

AIM:

EXPERIMENT-12-11

AIM- To identify the given inorganic salt $ZnCl_2$.

EXPERIMENT	OBSERVATIONS	INFERENCE
* PRIMARY TEST-		
1. Colour	White	Shows the absence of Cu^{2+} , Ni^{2+} , Co^{2+} .
2. Smell	No specific colour.	NH_4^+ , CH_3COO^- may be absent.
* DRY HEATING TEST		
3. Gas evolved	Colourless gas with pungent smell, white fumes with NH_3 presence	Cl^- may be present
4. Sublimate formed	No sublimation	NH_4^+ , I^- are absent.
5. Decrepitation	No decrepitation.	$Pb(NO_3)_2$, $NaCl$, KBr absent.
6. Residue	Yellow fns hot White fns cold	Zn^{2+} may be present.
7. Flame Test: Make salt + Conc. HCl paste	Shows a green flame	Zn^{2+} or Mn^{2+} may be present.
8. Dil H_2SO_4 test: Salt + Dil. H_2SO_4	No gas evolved	CO_3^{2-} , S^{2-} , NO_2^- , SO_3^{2-} may be absent.
9. $KMnO_4$ Test: Salt + dil H_2SO_4 and heat	When cold, Pink colour of $KMnO_4$ is discharged	Cl^- , Br^- , I^- may be present.

AIM:

10.	Conc. H_2SO_4 + Salt + Heat	Colourless gas, pungent smell, white fumes with NH_4 & white ppt of $AgNO_3$	Cl^- may be present.
11.	Heat a pinch of Salt and conc. $NaOH$	No NH_3 gas evolved	NH_4^+ absent.
12.	Shake the mix of Salt + Water	Solution is obtained	Label it as the original sol^n .
13.	To a part of 0.5 add 1-2 mL of dil HCl .	No ppt obtained	Group I absent (Pb^{2+} absent)
14.	Through the part of above, pass H_2S	No ppt obtained	Group II absent ($Pb^{2+}, Cu^{+2}, Ag^{2+}$, absent)
15.	To remaining sol^n . add pinch of NH_4Cl boil, cool, and add NH_4OH	No ppt formed	Group III absent (Fe^{3+}, Al^{3+} absent)
16.	To remaining sol^n add pinch of NH_4Cl pass H_2S gas.	white ppt obtained	Group IV present (Zn^{+2}, Mn^{+2} may be present).

RESULT -

Acidic Radical - Cl^- Basic Radical - Zn^{2+}

PRECAUTIONS -

- Don't heat wet test tube.

Expt. No. _____

Page No. 31

Date _____

AIM:

- ii) Don't inhale unidentified gases
- iii) Keep test-tube away from face while dry-heating.
- iv) Handle reagents carefully.

AIM:

EXPERIMENT - 12

AIM - To identify the given inorganic salt $ZnSO_4$.

EXPERIMENT	OBSERVATIONS	INFERENCE.
* PRIMARY TEST -		
1. Colour	white	Shows absence of Cu^{2+} , Fe^{3+} , Co^{2+} , Mn^{2+}
2. Smell	No specific smell	NH_4^+ , CH_3COO^- , S^{2-} may be absent.
* DRY HEATING TEST -		
3. Gas evolved	No gas evolved	SO_3^{2-} , Cl^- , Br^- , NO_3^- are absent.
4. Sublimate formed	No sublimation	NH_4^+ , I^- are absent.
5. Decoloration	Absent	$Pb(NO_3)_2$, $NaCl$, HBr are absent.
6. Residue	Hot \rightarrow Yellow Cold \rightarrow White	Zn^{2+} may be present.
7. Flame Test - Paste of Salt + Conc. HCl & heated	Green flame	Zn^{2+} or Mn^{2+} may be present.
8. Dil H_2SO_4 + Salt + Heat (if required)	No gas evolved	CO_3^{2-} , S^{2-} , NO_2^- , SO_3^{2-} may be absent.
9. Add $KMnO_4$ to Salt + dil H_2SO_4	Pink colour of $KMnO_4$ is discharged.	Cl^- , Br^- , I^- , $C_2O_4^{2-}$ may be absent.
10. Test for independent radical Boil a small amount.	A white ppt insoluble in conc. HCl obtained	SO_4^{2-} is present.

AIM:

of salt with dil. HCl,
filter it and add
few drops of BaCl_2
 Sol^n

11.	Heat a pinch of salt with conc. NaOH	No ammonia gas evolved.	NH_4^+ absent.
12.	To a part original solution add 1-2 ml of dil. HCl.	No ppt obtained	Group I absent (Pb^{2+} absent)
13.	Through a part of previous sol^n , pass H_2S gas	No ppt formed	Group II absent (Pb^{2+} absent)
14.	To remaining sol^n add pinch of NH_4Cl , boil the sol^n , cool it down and add excess NH_4OH	No ppt formed	Group III absent (Fe^{3+} , Al^{3+} absence)

RESULT -

Acid Radical - SO_4^{2-}

Basic Radical - Zn^{2+}

PRECAUTIONS -

- i) Never heat wet test-tube
- ii) Don't inhale unknown or poisonous gases.
- iii) Keep test-tube in a safe distance while dry-heating.
- iv) Handle reagents carefully.

AIM:

EXPERIMENT - 16.15

AIM - To identify the given inorganic salt $(\text{NH}_4)_3\text{PO}_4$.

EXPERIMENT	OBSERVATIONS	INFERENCE.
* PRIMARY TEST.		
1. Colour.	White	Shows absence of Ni^{2+} , Fe^{3+} , Co^{2+} , Mn^{2+} , NH_4^+ is present.
2. Smell	Ammoniacal smell	NH_4^+ is present.
* DRY HEATING TEST		
3. Gas evolved	Colourless gas, pungent and sweet smell, white fumes.	NH_4^+ may be present.
4. Sublimation	White sublimate	NH_4^+ may be present.
5. Swelling	Salt swells	PO_4^{3-} may be present.
6. Flame Test.		
Make a paste of salt + conc. HCl & perform flame test.	No specific smell	Ca^{2+} , Sr^{2+} , Ba^{2+} , Cu^{2+} , Zn^{2+} , Pb^{2+} are absent.
7. Dil H_2SO_4 treated with a pinch of salt and heat	No gas is evolved	CO_3^{2-} , SO_3^{2-} , S^{2-} , NO_2^- are absent.
8. To a pinch of salt add dil. H_2SO_4 (warm) & then add KMnO_4	Decolourise KMnO_4 from pink	Cl^- , Br^- , I^- , $\text{C}_2\text{O}_4^{2-}$ and Fe^{2+} are absent.

AIM:

9.	Conc. H_2SO_4 + Salt + Heat (if required)	No gas evolved	Cl^- , Br^- , I^- , NO_3^- , CH_3COO^- absent.
10.	Heat a pinch of Salt with conc. NaOH	Colourless gas, with Ammoniacal smell evolved.	Group zero is present (NH_4^+ Present)

RESULT-

Acid Radical - PO_4^{3-}

Basic Radical - NH_4^+

PRECAUTIONS-

- 1) Never heat a wet test tube.
- 2) Don't inhale gases which are unknown / poisonous.
- 3) Handle reagents very carefully.

AIM:

EXPERIMENT-17.16

AIM- To identify the given inorganic salt NH_4Br .

EXPERIMENT	OBSERVATIONS	INFERENCE
* PRIMARY TEST -		
1. Colour	white	Cu^{2+} , Ni^{2+} , Fe^{3+} , are absent.
2. Smell	Ammoniacal	NH_4^+ is present.
3. Gas evolved	Colourless gas, with characteristic smell gives white fumes when.	NH_4^+ may be present.
4. Sublimate formed	white sublimate	NH_4^+ may be present.
5. Flame Test: Make a paste of salt + conc. HCl. perform flame test.	No specific flame	Shows absence of Ca^{2+} , Ba^{2+} , Zn^{2+} , Pb^{2+} .
6. Treat a pinch of salt with dil H_2SO_4 & heat	No gas evolved	CO_3^{2-} , SO_3^{2-} , S^{2-} , NO_2^- , absent.
7. Salt + dil KMnO_4 heat it and add a drop of H_2SO_4	Pink colour of KMnO_4 decolourises	Cl^- , Br^- , I^- may be present.
8. Salt + conc. H_2SO_4 + heat (if required)	Reddish brown gas with pungent odour. Turns FeSO_4 into black	Br^- may be present.
9. Heat a pinch of salt with conc. NaOH	Colourless gas with ammoniacal smell	Group zero present (NH_4^+ present).

RESULT - 1) Acid Radical - Br^- 2) Basic Radical - NH_4^+

vision

Teacher's Signature: _____

Expt. No. 18.

Page No. 42

Date _____

PRECAUTIONS -

- 1) No heating of wet test tube.
- 2) Use test-tube away from body.
- 3) Handle reagents carefully.

AIM:

EXPERIMENT-18 17

AIM - To identify the given inorganic salt $(\text{NH}_4)_2\text{CO}_3$.

EXPERIMENT	OBSERVATIONS	INFERENCE.
1. Colour	White	Shows absence of Cu^{2+} , Fe^{2+} , Fe^{3+} , Co^{2+}
2. Smell	Ammoniacal	NH_4^+ is present.
3. Gas evolved	Colourless with characteristic smell gives white fumes when a Nessler's sol ⁿ brown	NH_4^+ maybe present.
4. Sublimation	white sublimate	NH_4^+ maybe is present.
5. Decapitation	no decapitation	$\text{Pb}(\text{NO}_3)_2$, NaCl is absent.
6. Flame Test Make paste of salt + conc. HCl & flame tested.	No specific flame	Ca^{2+} , Sr^{2+} , Ba^{2+} , Zn^{2+} , Pb^{2+} are absent.
7. Dil H_2SO_4 + salt + Δ	Colourless, odourless, gas with brisk effervescence	CO_3^{2-} may be present.
8. Salt + water	Salt does not dissolve	Involves CO_3^{2-} indicated
9. Salt + conc. H_2SO_4 + heat (if required)	No gas evolve	Cl^- , Br^- , I^- , NO_3^- , CH_3COO^- , $\text{C}_2\text{O}_4^{2-}$ are absent.
10. Salt + conc. H_2SO_4 + heat (if required)	No decolorisation	Indication of carbonate

RESULTS -

Acid Radical $\rightarrow \text{CO}_3^{2-}$ Basic Radical $\rightarrow \text{NH}_4^+$.

Expt. No. _____

Page No. 44

Date _____

AIM:

PRECAUTIONS -

- i) No heating of wet test-tubes.
- ii) Use test-tube away from body.
- iii) Handle reagents carefully.

AIM:

EXPERIMENT-20-19

AIM- To identify the given inorganic salt $Sr(NO_3)_2$

EXPERIMENT	OBSERVATION	INFERENCE.
1. Colour	white	shows absence of Ni^{2+} , Fe^{2+} , Fe^{3+} , Co^{2+} .
2. Smell	No specific smell	NH_4^+ , CH_3COO^- , S^{2-} are absent.
3. Gas evolved	NO_2 gas - Red-brown gas turns $FeSO_4$ into black.	NO_3^- may be present
4. Sublimate formed	No sublimation	$Pb(NO_3)_2$, $NaCl$, HBr is absent.
5. Decrepitation	No decrepitation	NH_4^+ , I^- are absent.
6. Residue	White residue which glows on heating	Ba^{2+} , Sr^{2+} , Ca^{2+} , Mg^{2+} may be present.
7. Flame Test: Salt + Conc. HCl & perform test	Crimson Red flame	Sr^{2+} may be present.
8. Salt + dil H_2SO_4 and heat sol ⁿ	No gas evolved	CO_3^{2-} , S^{3-} , NO_2^- , SO_3^{2-} is absent.
9. Salt + dil H_2SO_4 + Heat + few drops of $KMnO_4$	Pink colour of $KMnO_4$ was not discharged	Cl^- , Br^- , I^- , $C_2O_4^{2-}$, Fe^{2+} may be absent.
10. Heat a pinch of salt with conc. $NaOH$	No ammonia gas evolved	NH_4^+ absent.
11. Shake mix of Salt + water	Sol ⁿ is obtained	label as original Sol ⁿ .



Teacher's Signature : _____

Expt. No. _____

Date _____

AIM:

12.	To a part of 0.5 add 1-2 ml of dil HCl	No ppt obtained	Group I absent Pb^{2+} absent.
13.	Through a part of this sol ⁿ pass H_2S gas	No ppt formed	Group II absent
14.	Through a part of this sol ⁿ . pass H_2S gas.	No ppt formed	Group V present (Ca^{2+} , Ba^{2+} , Sr^{2+} may be present).

RESULT -

Acidic Radical - NO_3^- Basic Radical - Sr^{2+}

PRECAUTION -

- i) Don't heat wet test-tube.
- ii) Handle reagents carefully.
- iii) Don't inhale unknown gas.

AIM:

EXPERIMENT - 21 20

AIM: To identify functional group of aldehyde. $\left(\overset{\text{O}}{\parallel} \text{C} - \text{H} \right)$

EXPERIMENT	OBSERVATION	INFERENCE
1. Test for unsaturation	Brown color of bromine not discharged	No Unsaturation is present.
2. Test for Carboxylic group	No effervescence	Carboxylic group is absent
3. Test for phenolic group	No green or violet colour obtained	Phenolic group is absent.
4. Test for alcoholic group	No effervescence	Alcoholic group is absent.
5. Test for Carbonyl group	Orange - yellow ppt formed	Carbonyl group is present - may be an aldehyde or a ketone.
6. Test for Carbonyl group	Silver mirror formed on inner side of test-tube	Aldehyde is present.
7. Test for Amine To a small amount of organic liq in test-tube add 1-ml conc. HCl & CHCl ₃ . Also add 2ml of alc. KOH + Heat.	No offensive smelling gas is evolved	Amino group absent.

RESULT -

The set of tests prove the presence of $\left(\overset{\text{O}}{\parallel} \text{C} - \text{H} \right)$ aldehyde functional group.

Expt. No. _____

Date _____

AIM:

PRECAUTIONS -

- i) use freshly prepared solutions.
- ii) Keep a safe distance from test-tube while heating
- iii) Avoid inhalation of any fumes evolved which are unknown.
- iv) Use a lab coat & gloves while dealing with corrosive chemicals.

AIM:

EXPERIMENT-22 21

 AIM - To identify functional group of ketone. $\left(- \overset{\text{O}}{\parallel}{\text{C}} - \right)$

EXPERIMENT	OBSERVATION	INFERENCE
1. Test for unsaturation dissolve 0.2 ml of CCl_4 then add Br_2 water	Brown colour of Bromine not discharged.	No unsaturation is present.
2. Test for Carboxyl group 0.2 ml of Compound + Pinch of NaHCO_3	No effervescence	Carboxylic acid, group is absent.
3. Test for phenolic group - 0.2 ml Compound + 2-3 ml of neutral FeCl_3 solution.	No green or violet colour obtained	Phenolic group is absent.
4. Test of alcoholic group - Small piece of sodium + 1 ml of given compound	No effervescence	Alcoholic group is absent.
5. Test for Carbonyl group, shake 0.2 ml of 2,3-di-nitro phenyl hydrazine	Orange - Yellow ppt formed.	Carbonyl group is present (Aldehyde / ketone).

AIM:	<p>6. Test for Ketonic group - 0.5 ml Compound + 0.1 ml m-dinitro benzene + 1 ml dil. NaOH.</p>	<p>Violet colour obtained that slowly fades.</p>	<p>Ketonic group is present.</p>
7.	<p>Confirmatory for Ketone - Dissolve a crystal of Sodium nitro-prusside in distilled water + 0.5 g/ml of Compound + NaOH drop-wise.</p>	<p>Red colour is obtained.</p>	<p>Ketone is confirmed.</p> $\begin{array}{l} \text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_3 + \text{OH}^- \\ \text{CH}_3 - \underset{\text{O}}{\underset{ }{\text{C}}} - \text{CH}_3 + \text{H}_2\text{O} \end{array}$

RESULT -

The given organic compound contains Ketone group $(-\overset{\text{O}}{\parallel}{\text{C}}-)$

PRECAUTIONS -

- FeCl_3 solⁿ should be freshly prepared.
- Br_2 water should be handled carefully.
- Unreacted Na metal should not be thrown in sink directly.

AIM:

EXPERIMENT-23-22

AIM- To identify the functional group of Alcohol (-OH).

EXPERIMENT	OBSERVATIONS	INFERENCE
1. Test for unsaturated dissolve 0.2 ml of Compound. In 2 ml of CCl_4 then add Br_2 water.	Brown colour of Bromine not dischar- ged	No unsaturation present.
2. Test for the Carboxylic group - 0.2 ml Compound + pinch of $NaHCO_3$	No effervescence	Carboxylic group is absent.
3. Test for phenolic group - 0.2 ml organic Compound + 2-3 ml $FeCl_3$ sol^n	No green or violet colour obtained.	Phenolic group is absent.
4. Test for carbonyl group: shake 0.2 ml of the Compound + 2-3 ml of 2,3 di-nitro phenyl hydrozine.	No ppt obtained	Carbonyl group is absent.

VISION

Teacher's Signature _____

Expt. No. _____

AIM:

5. Test for alcoholic group: Small piece of Na + 1ml of compound

Effervescence obtained

Alcohol group is present.

RESULT-

The given organic compound contains alcoholic ($-\text{OH}$) group.

PRECAUTIONS-

- i) FeCl_3 solⁿ should be freshly prepared
- ii) Br_2 water should be handled carefully.
- iii) Unreacted ' Na ' should not be disposed directly into the sink.

Identify the functional present in the given organic compound (Carboxylic acid)

Experiment	Observation	Inference
Test for Unsaturation Dissolve 0.2 ml Comp in 2 ml CCl ₄ then add Br ₂ water	Brown colour of bromine not discharged	No Unsaturation is present.
Test for phenolic group Add 0.2 ml compound 2-3 ml neutral FeCl ₃ sol ⁿ	No green violet colour obtained	phenolic group is present.
Test for alcoholic group Small piece of Na ⁺ 1 ml of given liq	no effervescence	Alcoholic group is absent.
Test for Carbonyl group -Shook 0.2 ml of Comp with 2-3 ml of 2,3,5-dinitrophenyl hydrazine	no orange yellow ppt formed	Carbonyl groups Aldehyde and Ketone are absent
Test for Carboxylic acid group -0.2 ml of Comp + pinch of NaHCO ₃	effervescence obtained	Carboxylic group is present

Confirmation test for COOH grp - 0.1g Comp + 1ml of ethyl alcohol and 1-2 drop of conc. H_2SO_4 + heat the mix ⁿ mixture on a beaker & Condensing Water	A faintly smell obtained	$-\text{COOH}$ is Confirmed $\text{RCOOH} + \text{C}_2\text{H}_5\text{OH}$ $\xrightarrow{\text{H}_2\text{SO}_4}$ RCOOC_2H_5 $+ \text{H}_2\text{O}$ faintly smell
--	--------------------------------	--

Result \rightarrow The organic compound contains
 Carboxylic ($-\text{COOH}$) group.

precautions \rightarrow

- FeCl_3 Solⁿ should be freshly prepared.
- Br_2 water should be handled carefully.
- Unreacted Na metal should not be thrown
 in sink directly.